



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

1; diseases of the nervous system, 1; of the circulatory system, 5; of the respiratory system, 3; of the digestive system, 3; of the urinary system, 1; congenital debility, 1; violence, 4; diseases badly defined, none; poisoning, 1. Natives, 17; foreigners, 7. Daily average of deaths, 3.41.

*Mortality of Rio de Janeiro for the week ended May 14, 1905.*

During the week ended on the above-named date there were in all 281 deaths. Of this number 18 were caused by yellow fever, with 42 new cases reported, of which number 29 were confirmed before the issuance of the report. Nine of the deaths occurred in the hospital São Sebastião, and the remainder in 6 of the city districts. At the close of the week there were 41 cases in the hospital and 3 other cases under observation.

This report does not seem so favorable as the last one, and it is very difficult, indeed impossible, to predict the future from the information at hand.

Variola caused only 3 deaths with 8 new cases, and at the end of the week there were 24 cases in the hospital São Sebastião.

There were no cases and no deaths from bubonic plague, and there is but 1 case remaining in the hospital, convalescent.

There have been no cases of plague in this city for 3 weeks, and there has been only 1 death during that period.

Other causes of death were: Measles, 1; whooping cough, 1; grippe, 11; enteric fever, 1; beriberi, 1; erysipelas, 1; malarial fevers, 2; tuberculosis, pulmonary form, 58; other forms 3; septicæmia, 1; cancer, 2; diseases of the nervous system, 25; of the circulatory system, 41; of the respiratory system, 37; of the digestive system, 40; of the urinary system, 10; puerperal septicæmia, 2; diseases of the skin, 1; congenital debility, 9; senile debility, 5; violence, 7; and suicide, 1.

By localities: In houses, homes, hotels, etc., 187; in civil hospitals, 27; in military hospitals, 3; in Santa Casa de Misericórdia, 56; in asylums, convents, etc., 5; in localities unknown, 3.

Natives, 201; foreigners, 78; and nationality unknown, 2. All yellow fever deaths this week were among foreigners.

Daily average of deaths, 40.14; same for the preceding week, 36.14; and for the corresponding week of 1904, 49.85.

Rate per each 1,000 of the population, estimated at 905,000, 16.19.

Highest range of the thermometer, centigrade, 27°; lowest, 17.3°; average for the week, 21.29°. Total rainfall for the week, 5.16 mm.

*Recurrence of yellow fever at Rio de Janeiro following on commencement of harbor works—Mortality among fish in bay.*

The following report from Consul-General Seeger, dated May 20, has been transmitted to this Bureau from the Department of State:

Yellow fever, which has prevailed in this port for many years, had been almost completely eradicated, when recently the following coincidences were observed in the Rio port: Commencement of dredging in an infected part of the harbor for the new harbor works; great mortality among the fish of the bay; recurrence of the yellow fever in an epidemic form, and this in the winter season.